

# • PRINTER RUSH • (PTO ASSISTANCE)

		<del></del>	<u>Ulm</u>	GAU:	1646						
From:	$e\omega c$	Location:	(ibc) FMF FDC	Date:	1 1/8						
I F W Tracking #: 06018431 Week Date: 10.4.04											
	DOC CODE  1449  IDS CLM IIFW SRFW DRW OATH 312 SPEC	DOC DATE	MISCELL Continuing Foreign Price Document I Fees Other	Data ority							
[RUSH] MESSAGE:  - First line of continuing data on Palm sheet is  not in specification.  - "CIP" is cited for 2nd and 3rd times on  bib sheet, but only claims priority is  stated in specification.  Please advisa  Thank you  Ew C											
[XRUSH] RES	SPONSE:										
				INITI	ALS:						

NOTE: This form will be included as part of the official USPTO record, with the Response document coded as XRUSH.

REV 10/04

30

35

10

# NOVEL MEMBERS OF THE CAPSAICIN/VANILLOID RECEPTOR FAMILY OF PROTEINS AND USES THEREOF

## 5 Related Applications

This application claims priority to U.S. provisional Application No. 60/108,322, filed on November 13, 1998, U.S. provisional Application No. 60/114,078 filed on December 28, 1998, U.S. Patent Application Serial No.: 09/258,633 filed on February 26, 1999, and U.S. Patent Application Serial No.: 09/421,134 filed on October 19, 1999, incorporated herein in their entirety by this reference.

#### Background of the Invention

Pain is initiated when the peripheral terminals of a subgroup of sensory neurons are activated by noxious chemical, mechanical or thermal stimuli. These neurons, called nociceptors, transmit information regarding tissue damage to pain-processing centres in the spinal chord and brain (Fields, H.L. Pain, McGraw-Hill, New York, 1987). Nociceptors are characterized in part, by their sensitivity to capsaicin, a vanilloid-containing compound, and a natural product of capsicum peppers that is the active ingredient of many "hot" and spicy foods. In mammals, exposure of nociceptor terminals to capsaicin leads initially to excitation of the neuron and the consequent perception of pain and local release of inflammatory mediators. With prolonged exposure, nociceptor terminals become insensitive to capsaicin, as well as to other noxious stimuli (Szolcsanyi, J. in Capsaicin in the Study of Pain (ed. Wood, J.) 1-26 (Academic, London, 1993). This latter phenomenon of nociceptor desensitization underlies the seemingly paradoxical use of capsaicin as an analgesic agent in the treatment of painful disorders ranging from viral and diabetic neuropathies to rheumatoid arthritis (Campbell, E. in Capsaicin and the Studyof Pain (ed. Wood, J.) 255-272 (Academic, London, 1993); Szallasi, A. et al. (1996) Pain 68, 195-208). Some of this decreased sensitivity to noxious stimuli may result from reversible changes in the nociceptor, but the long-term loss of responsiveness can be explained by death of the nociceptor or destruction of its peripheral terminals following exposure to capsaicin (Jancso, G. et al. (1977) Nature 270, 741-743).

The cellular specificity of capsaicin action and its ability to evoke the sensation of burning pain have led to speculation that the target of capsaicin action plays an important physiological role in the detection of painful stimuli. Indeed, capsaicin may elicit the perception of pain by mimicking the actions of a physiological stimulus or an endogenous ligand produced during tissue injury (James, I.F., Kinkina, N.N. & Wood, J.N. in Capsaicin in the Study of Pain (ed. Wood, J.N.) 83-104 (Academic, London, 1993).

Caterina M.J. et al. have recently determined the molecular basis underlying this phenomenon by characterizing a functional cDNA that encodes a vanilloid receptor (VR-1) in rat



## United States Patent and Trademark Office

COMMISSIONER FOR PATENTS
UNITED STATES PATENT AND TRADEMARK OFFICE
WASHINGTON, D.C. 20231

# 

**Bib Data Sheet** 

SERIAL NUMBER 09/587,111	FILING DATE 06/02/2000 RULE	(	CLASS 435	GROUP ART UNIT		ATTORNEY DOCKET NO. MNI-062CP2DV1					
APPLICANTS Rory A.J. Curtis, Southborough, MA;  ** CONTINUING DATA **********************************											
Foreign Priority claimed 35 USC 119 (a-d) condition met Verified and Acknowledged  ADDRESS	itiats	STATE OR COUNTRY MA			TOT/ CLAII 50	MS	INDEPENDENT CLAIMS 6				
959 TITLE  Novel members of the capsaicin/vanilloid receptor family of proteins and uses thereof											
RECEIVED No.	S: Authority has been given in Paper to charge/credit DEPOSIT ACCOUNT for following:				All Fees  1.16 Fees (Filing)  1.17 Fees (Processing Ext. of time)  1.18 Fees (Issue)  Other  Credit						